



P16-31. Skewed HIV-1-Specific CD4+ Th2 Helper Cell Contribution in Progressive HIV-1 Infection

Citation

Chevalier, M., A. Pyo, J.S. Jolin, M. Addo, D.S. Kwon, I. Toth, B. Walker, and H. Streeck. 2009. P16-31. Skewed HIV-1-specific CD4+ Th2 helper cell contribution in progressive HIV-1 infection. Abstracts from AIDS Vaccine 2009 poster and podium presentations. *Retrovirology* 6(Suppl 3): P260.

Published Version

doi:10.1186/1742-4690-6-S3-P260

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:5978767>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Poster presentation

Open Access

P16-31. Skewed HIV-1-specific CD4⁺ Th2 helper cell contribution in progressive HIV-1 infection

M Chevalier*, A Pyo, JS Jolin, M Addo, DS Kwon, I Toth, B Walker and H Streeck

Address: Ragon Institute of MGH, MIT and Harvard, Boston, MA, USA

* Corresponding author

from AIDS Vaccine 2009
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, **6**(Suppl 3):P260 doi:10.1186/1742-4690-6-S3-P260

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P260>

© 2009 Chevalier et al; licensee BioMed Central Ltd.

Background

CD4⁺ T helper cells play a crucial role in the orchestration of the immune system. While CD4⁺ Th1 responses provide important helper signals to the survival and maturation of CD8⁺ T cells, CD4⁺ Th2 responses are involved in B cell maturation and antibody class switching. However, very little is known about the role and contribution of HIV-1-specific CD4⁺ T helper responses to the control of viral replication. Here we show that the presence of Gag-specific CD4⁺ Th1 responses and Th2 responses against Env are associated with viral control, while changes in this pattern are associated with disease progression.

Methods

We cross-sectionally screened 21 subjects with chronic HIV-1 infection (>10,000 copies/mL) and 15 elite controllers (<50 copies/mL) for HIV-1-specific CD4⁺ T helper responses by flow cytometry before and after cultivation in helper subset polarizing conditions (Th1: IFN γ /IL12; Th2: IL4/anti-IFN γ ; Th17: anti-IL4/anti-IFN γ /TGF β /IL6).

Results

HIV-1-specific CD4⁺ T helper responses were detected in similar frequencies in chronic HIV-1 infection and elite controllers. However, we observed in HIV-1 elite controllers significantly more CD4⁺ T cell responses consisting of a Th1 (IFN γ /IL2⁺) CD4⁺ T cell phenotype. These responses were predominantly directed against the HIV-1 Gag protein. In contrast, in subjects with chronic-progressive HIV-1 infection Th1 responses were directed against

envelope in a higher frequency ($p < 0.02$). Interestingly, while the HIV-1-specific Th2 responses in elite controllers were preferentially directed against envelope, we observed a skewed over-representation of Th2 responses against Gag in subjects with chronic-progressive HIV-1 infection. These responses were also associated with higher activation levels.

Conclusion

Our data suggest that the presence of a distinct CD4⁺ T helper cell pattern is associated with viral control. The presence of Gag-specific Th1 responses and Env-specific Th2 responses suggests a potential important role for the helper signals for B and T cell responses. This finding might be highly important for vaccine design.